

ELEMENTARY DERIVATION OF THE CHIRAL ANOMALY

EDWIN LANGMANN AND JOUKO MICKELSSON

Theoretical Physics, Royal Institute of Technology, S-10044, Stockholm, Sweden

December 15, 1994

ABSTRACT An elementary derivation of the chiral gauge anomaly in all even dimensions is given in terms of noncommutative traces of pseudo-differential operators.

The ordinary trace functional acting on $N \times N$ matrices is always defined and has the fundamental property that $\text{tr}[A, B] = 0$. In infinite dimensions the concept of trace becomes more subtle. Even for bounded Hilbert space operators the trace does not usually make sense (the sum of diagonal elements either diverges or depends on the bases). In the case of pseudo-differential operators there exist alternative definitions. An important trace is given by the operator residue, $[W]$, which picks up precisely the logarithmically diverging part of the ordinary trace and gives it a finite value; for a suitable class of operators this is equal to the Dixmier trace used extensively in noncommutative geometry, $[C]$.

In certain cases important in applications of operator theory to quantum field theory there is an alternative way to make sense of the logarithmically diverging piece. Let d be the dimension of the underlying space-time manifold. It may happen that even though the trace of a PSDO is formally diverging (by power counting, the logarithmically diverging borderline case being degree $= -d$) it has a finite value as a limit $\Lambda \rightarrow \infty$ of the ultraviolet cut-off trace tr_Λ . This is the case when the leading symbol of the PSDO is a total derivative in momentum space; in that case the cut-off trace becomes a finite surface integral (plus possibly a finite ordinary trace of

the lower order contributions). The most notable example for this behavior is when the PSDO can be written as a commutator. If the degree (in momentum space) of the commutator is precisely $-d$ then the trace depends only on the leading symbol and is given exactly by a surface integral. This fact has been used in calculations involving Schwinger terms in quantum field theory, [M],[LM],[CFNW],[L]. In this note we shall show that the symbol calculus of PSDO's combined with the use of the noncommutative trace, as discussed above, gives a simple derivation of the known formula for the chiral anomaly (see the reviews in [R] and [TJZW] for earlier approaches to this problem or [BGV] for a more mathematical discussion of the subject).

Although the use of symbol calculus of pseudodifferential operators has always been implicit in calculations of Feynman diagrams we find that it is of great advantage to use the symbol calculus in a systematic way. We believe that the theory is not yet fully exploited in quantum physics and more applications are waiting, in addition to those in [M],[LM],[CFNW],[L].

Let $D_A = \gamma^\mu(-i\partial_\mu + A_\mu P_+)$ be the *chiral* (nonselfadjoint) Dirac operator in $\mathbb{R}^d$, $d = 2n$ even. Here A is a Yang-Mills vector potential with values in a Lie algebra $\mathfrak{g}$ of a compact Lie group G and $P_+ = \frac{1}{2}(1 + \gamma_{d+1})$ is the chiral projection operator with $\gamma_{d+1} = (-i)^n \gamma_1 \gamma_2 \dots \gamma_{2n}$. The $2^n \times 2^n$ gamma matrices satisfy the euclidean anticommutation relations $\gamma_i \gamma_j + \gamma_j \gamma_i = 2\delta_{ij}$. The Lie algebra $\mathfrak{g}$ acts on the components of the Dirac field ψ through a finite-dimensional unitary representation ρ of G . Thus ψ takes values in $V = \mathbb{C}^{2^n} \otimes \mathbb{C}^N$, where N is the dimension of ρ .

We shall apply the symbol calculus of pseudo-differential operators and therefore we assume that the potential A and the gauge transformations have compact support.

A PSDO A is given by its symbol $a(x, p) = \sigma(A)(x, p)$ which is a smooth function of the coordinates $x \in \mathbb{R}^d$ and momenta $p \in \mathbb{R}^d$, [H]. We shall consider *the restricted class* of PSDO's which admit an asymptotic expansion of the symbol as

$$a(x, p) \sim a_k(x, p) + a_{k-1}(x, p) + a_{k-2}(x, p) + \dots$$

where k is an integer and each a_j is a homogeneous matrix valued function of the momenta of degree j with $|a_j| \sim |p|^j$ as $\sqrt{x^2 + p^2} \rightarrow \infty$. The

asymptotic expansion for the product of two PSDO's is given by the formula

$$(1) \quad a * b \sim \sum \frac{(-i)^{|m|}}{m!} [(\partial_p)^m a(x, p)] [(\partial_x)^m b(x, p)],$$

where the sum is over all sets of nonnegative integers $m = (m_1, \dots, m_d)$, $|m| = m_1 + \dots + m_d$, $\partial_x^m = (\frac{\partial}{\partial x_1})^{m_1} \dots (\frac{\partial}{\partial x_d})^{m_d}$, etc., and $m! = m_1! \dots m_d!$.

The trace (if it exists) of a matrix valued PSDO A is given as

$$(2) \quad \text{tr } A = \frac{1}{(2\pi)^d} \int \text{tr } a(x, p) dx dp.$$

The trace exists if the symbol $a(x, p)$ behaves asymptotically $|p|^{-d-\epsilon}$ for a positive ϵ , assuming that $a(x, p)$ has compact support in x (this condition is satisfied in the cases of interest to us). The borderline case $\epsilon = 0$ is important to us: in that case the trace diverges logarithmically *except when the leading symbol $a_{-d}(x, p)$ is a total derivative in momentum space*. If it is a total derivative the trace can be written as a (finite) surface integral of some symbol which is of degree $-2n+1$. For the restricted class of PSDO's A with a symbol a of degree k we can expand the *cut-off trace* $\text{tr}_\Lambda A$ in powers of the momentum space cut-off Λ ,

$$(3) \quad \begin{aligned} \text{tr}_\Lambda A &= \frac{1}{(2\pi)^d} \int_{|p| \leq \Lambda} dp \int dx \text{tr } a(x, p) \\ &= c_{k+2n} \Lambda^{k+2n} + c_{k+d-1} \Lambda^{k+d-1} + \dots c_{\log} \log \Lambda + c_0 \Lambda^0 + c_{-1} \Lambda^{-1} + \dots \end{aligned}$$

The logarithmic term arises from the trace of the degree $-d$ piece in the symbol and $c_{\log}(A)$ is proportional to the operator residue of A . The regularized trace of a PSDO is then defined as

$$\text{TR } A = c_0(A)$$

in the above expansion. If $\deg A < -d$ then the regularized trace is equal to the ordinary trace $\text{tr } A$.

The degree of $a * b$ is equal to the sum of degrees of a and b . The leading term in $a * b$ is just the matrix product ab of the symbols. If $\deg a + \deg b = p$ the highest order term which contributes to the trace $\text{tr } [A, B]$ is of degree $p - 1$ since the finite-dimensional commutator traces in V vanish identically.

Our calculation of the anomaly is based on the fact that we can write the gauge variation of the effective action as a sum of traces of commutators. For bounded operators A, B the trace $\text{tr } [A, B]$ is stable with respect to perturbations of A, B by

trace class operators. Therefore, it depends only on the asymptotic expansion of the symbols of A and B . This is because two PSDO's with the same asymptotic expansion can differ only by a so-called infinitely smoothing operator. An infinitely smoothing operator is always trace-class (for compactly supported symbols in x space).

The effective action for chiral fermions coupled to external Yang-Mills field is defined as (the logarithm of) $\det_{reg}(D_A)$ where 'reg' means that some sort of regularization is needed for the convergence of the infinite-dimensional determinant. We shall use the following regularization. Formally, the logarithm of a determinant is equal to the trace of the logarithm of the operator. Thus we define tentatively

$$\log \det_{reg}(D_A) = \text{TR} \log(D_A).$$

Of course, the right-hand-side is ill-defined when D_A is not invertible and for invertible operators one has to fix the phase of the logarithm somehow. It will turn out that for the computation of chiral anomaly the form of the 'infra-red' regularization of the Dirac operator is inessential (since the anomaly will be a surface integral in momentum space). In the following discussion, leading to the definition (5) of the effective action, we shall therefore not be specific about the choice of the infrared regularization; we shall return to this problem later.

The operator $\log(D_A)$ is not a classical pseudo-differential operator, but we can write $\log(D_A) = \log(D_0) + W_A$, where W_A is a classical pseudodifferential operator of degree zero. We can then drop an (infinite) constant from the definition of the effective action by declaring it to be equal to $-\text{TR} W_A$. The operator W_A is very complicated function of A because

$$\log(ST) = \log(S) + \log(T) + F(S, T)$$

where $F(S, T) = \frac{1}{2}[\log(S), \log(T)] +$ higher commutators. When computing the regularized trace one is not allowed to drop the commutator terms because in general $\text{TR}[S, T] \neq 0$. However, only a finite number of commutators are relevant, giving a contribution to the effective action which is a finite differential polynomial. This is seen as follows. In our case $S = D_0$ and $T = D_0^{-1}D_A = 1 + D_0^{-1}\mathcal{A}_+$ where $\mathcal{A}_+ = \gamma^\mu A_\mu P_+$. Thus

$$(4) \quad \log(T) = D_0^{-1}\mathcal{A}_+ - \frac{1}{2}(D_0^{-1}\mathcal{A}_+)^2 + \frac{1}{6}(D_0^{-1}\mathcal{A}_+)^3 + \dots$$

It follows that there are only a finite number of terms in the expansion of the logarithm which are of degree $> -d$ and which can contribute to the trace of the commutator. Furthermore, for each term we need to compute derivatives in momentum space (for the asymptotic expansion (1)) up to a finite degree, because each derivative decreases the degree by one unit. Finally, in the multiple commutator expansion for $F(S, T)$ we need to take only a finite number of commutators with T because the degree of $\log(T)$ is -1 .

The conclusion is that

$$\text{TR } W_A = \text{TR} \log(1 + D_0^{-1} \mathcal{A}_+) + P(A)$$

where $P(A)$ is a finite differential polynomial in A . After performing this polynomial renormalization we define

$$(5) \quad S_{eff}(A) = \text{TR} \log(1 + D_0^{-1} \mathcal{A}_+).$$

This definition gives also a resolution to the phase problem for the logarithm. For small potentials A , after fixing an infrared regularization for D_0 (e.g. $D_0^{-1} \rightarrow (D_0 + i\epsilon)^{-1}$) one can expand $\log(1 + D_0^{-1} \mathcal{A}_+)$ as a convergent power series in $D_0^{-1} \mathcal{A}_+$. A similar expansion can be performed around an arbitrary background potential B by replacing $D_0 \rightarrow D_B$.

The chiral anomaly is defined as $\omega(X; A) = \mathcal{L}_X S_{eff}(A)$, where $\mathcal{L}_X$ denotes the Lie derivative in the direction of an infinitesimal gauge transformation $X : \mathbb{R}^{2n} \rightarrow \mathbf{g}$, $(\mathcal{L}_X f)(A) = \frac{d}{dt} f(-ie^{-itX} de^{itX} + e^{-itX} A e^{itX})|_{t=0}$. We shall show below that the anomaly can be written as a regularized trace of commutators of PSDO's which then leads automatically to a finite differential polynomial expression for the anomaly.

Using $-i\mathcal{L}_X \mathcal{A} = -i\mathcal{A}X + [\mathcal{A}, X]$, where $\mathcal{A}X = \gamma^\mu \partial_\mu X$, $\mathcal{A} = \gamma^\mu A_\mu$, etc., we get a symbol formula for $\omega(X; A)$,

$$(6) \quad \omega(X; A) = \sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{k} \sum_{j=0}^{k-1} \text{TR} \left(\frac{1}{D_0} \mathcal{A}_+ \right)^j \left\{ \frac{1}{D_0} \mathcal{A}X P_+ + \frac{1}{D_0} i[\mathcal{A}, X] P_+ \right\} \left(\frac{1}{D_0} \mathcal{A}_+ \right)^{k-j-1}.$$

(In computing the traces a fixed infrared regularization $1/(D_0 + i\epsilon) \rightarrow 1/(D_0 + i\epsilon)$ is

used.) On the other hand,

$$(7) \quad \begin{aligned} \text{TR} [\log(1 + \frac{1}{D_0} \mathcal{A}_+), iX] &= \sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{k} \sum_{j=0}^{k-1} \\ \text{TR} (\frac{1}{D_0} \mathcal{A}_+)^j &\left(-\frac{1}{D_0} \not{D} X \frac{1}{D_0} \mathcal{A}_+ + \frac{i}{D_0} [\mathcal{A}, X] P_+ \right) (\frac{1}{D_0} \mathcal{A}_+)^{k-j-1}. \end{aligned}$$

Disregarding commutator terms this expression is the same as the above formula for $\omega(X; A)$. More precisely,

$$(8) \quad \begin{aligned} \omega(X; A) &= \text{TR} [\log(1 + \frac{1}{D_0} \mathcal{A}), iX P_+] \\ &+ \sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{k(k+1)} \sum_{j=0}^{k-1} \text{TR} \left[(\frac{1}{D_0} \mathcal{A})^j (\frac{1}{D_0} \not{D} X), (\frac{1}{D_0} \mathcal{A})^{k-j} P_+ \right]. \end{aligned}$$

We have used the projection operator property $P_+^2 = P_+$ and the fact that P_+ commutes with a product of a pair of γ matrices. Since the anomaly is a sum of traces of commutators it really depends only on the asymptotic expansion of the operators involved.

Actually, there is a delicate point in the above argument which shows the difference between the terms arising from the γ_{d+1} part (the nontrivial part of the anomaly) or from '1' in the projection $P_+ = \frac{1}{2}(1 + \gamma_{d+1})$. Remember that we have to make an infrared regularization $\frac{1}{D_0} \rightarrow \frac{1}{D_0 + i\epsilon}$ in order to make sense of the traces. But the regularized operator $\frac{1}{D_0 + i\epsilon} \mathcal{A}$ does not commute with γ_{d+1} which prevents us of writing the gauge variation of the operator $\log(1 + \frac{1}{D_0 + i\epsilon} \mathcal{A} P_+)$ exactly as a sum of commutators; but this does not have an effect on the computation involving traces. The reason is that the nonzero commutators $[P_+, \frac{1}{D_0 + i\epsilon} \mathcal{A}]$ affect only terms arising from the commutator $[\log(1 + \frac{1}{D_0 + i\epsilon} \mathcal{A} P_+), iX]$. The ultraviolet regularized trace of the commutator depends only on the degree $-d$ part of the asymptotic expansion of the commutator. Thus the choice of the infra-red regularization does not have any effect on the result and can be ignored (see the explicit computations below); the remaining traces will be surface integrals in the momentum space.

A basic formula for the γ -matrix algebra which we need several times is

$$(9) \quad \text{tr } \gamma_{i_1} \gamma_{i_2} \cdots \gamma_{i_p} \gamma_{2n+1} = \begin{cases} 0, & \text{if } p < 2n \\ (2i)^n \epsilon_{i_1 i_2 \dots i_{2n}}, & \text{if } p = 2n. \end{cases}$$

We shall also use $\pi(D_0)(x, x) = \not{D} - \not{x} \not{\partial} x$

The case $d = 2$. Since now $\text{TR}[A, B] = 0$ when $\deg(A) + \deg(B) \geq 2$, the only nonvanishing term is

$$(10) \quad \omega(X; A) = \text{TR} \left[\frac{1}{D_0} \not{A}, iX P_+ \right].$$

The form of the infrared regularization at $p = 0$ is irrelevant, since the trace of the commutator will be a integral of a total derivative in momentum space.

$$\begin{aligned} \omega(X; A) &= \frac{1}{(2\pi)^2} \int dx \int dp (-i) \text{tr} \frac{\partial}{\partial p_\mu} \left(\frac{1}{\not{p}} \right) \not{A} \frac{\partial}{\partial x_\mu} iX P_+ \\ &= \frac{1}{(2\pi)^2} \int dx \int_{|p|=\infty} dp \text{tr} \frac{p_\mu}{|p|} \frac{\not{p}}{|p|^2} \not{A} \partial_\mu X P_+ \\ &= \frac{1}{4\pi} \int dx \text{tr} \gamma^\mu \gamma^\nu A_\nu \partial_\mu X \frac{1}{2} (1 + \gamma_3) \\ (11) \quad &= \frac{i}{4\pi} \int dx \epsilon^{\mu\nu} \text{tr} A_\mu \partial_\nu X + \frac{1}{4\pi} \int dx \text{tr} A_\mu \partial_\mu X. \end{aligned}$$

The real part (the second term) is a coboundary of the local functional $f(A) = \frac{1}{8\pi} \int dx \text{tr} |A|^2$ and thus a trivial 1-cocycle. Actually, the real part must be a trivial cocycle in any space-time dimension on cohomological grounds. Note also that the real part is coming from the term 1 in $P_+ = \frac{1}{2}(1 + \gamma_{2n+1})$. This term is the gauge anomaly for the nonchiral determinant $\frac{1}{2} \text{TR} \log(1 + \frac{1}{D_0} \not{A})$. We shall show later explicitly that really $\mathcal{L}_X \text{TR} \log(1 + \frac{1}{D_0 + i\epsilon} \not{A}) = \mathcal{L}_X P_d(A)$ where P_d is a local differential polynomial in A in any dimension d .

The case $d = 4$. Now the relevant terms contributing to the commutator trace are

$$\begin{aligned} \omega(X; A) &= -\frac{1}{2} \text{TR} \left[\left(\frac{1}{D_0} \not{A} \right)^2, iX P_+ \right] + \frac{1}{3} \text{TR} \left[\left(\frac{1}{D_0} \not{A} \right)^3, iX P_+ \right] \\ &\quad - \frac{1}{6} \text{TR} \left[\frac{1}{D_0} \not{A} \frac{1}{D_0} \not{A} X, \frac{1}{D_0} \not{A} P_+ \right] - \frac{1}{6} \text{TR} \left[\frac{1}{D_0} \not{A} X, \left(\frac{1}{D_0} \not{A} \right)^2 P_+ \right]. \end{aligned}$$

The actual computation of the various commutators and traces is a straight-forward application of (1) for the product of PSDO's and the γ matrix algebra (9). The result for the imaginary part is

$$(12) \quad \omega_\gamma(X; A) = \frac{-1}{24\pi^2} \int \text{tr} dX (-i dA A + \frac{1}{2} A^3)$$

where the exterior algebra notation has been used for the integrand. The real part is again trivial and can be written as a coboundary of a suitable differential polynomial in A .

The general case

First we regroup the terms in (8). We can write

$$(13) \quad -i\omega(X; A) = \sum_{m=1}^{\infty} \frac{(-1)^{m-1}}{m(m+1)} \sum_{j=0}^m \text{TR} \left\{ \left[\left(\frac{1}{D_0} A \right)^j, X \left(\frac{1}{D_0} A \right)^{m-j} P_+ \right] - \right. \\ \left. - \left[\left(\frac{1}{D_0} A \right)^j X, \left(\frac{1}{D_0} A \right)^{m-j} P_+ \right] + \left[\left(\frac{1}{D_0} A \right)^j \frac{1}{D_0} [D_0, X], \left(\frac{1}{D_0} A \right)^{m-j} P_+ \right] \right\}.$$

The right-hand side can be written in a more compact form, involving the inverse of the Dirac operator D_A , by the following trick. Using the identity

$$\frac{-1}{m(m+1)} = \int_0^1 t^m \left(1 - \frac{1}{t}\right) dt$$

we get

$$(14) \quad \omega(X; A) = \int_0^1 \nu(X; tA) \frac{t-1}{t} dt$$

with

$$(15) \quad -i\nu(X; A) = \sum_{m=0}^{\infty} \sum_{j=0}^m (-1)^m \text{TR} \left\{ \left[\left(\frac{1}{D_0} A \right)^j, X \left(\frac{1}{D_0} A \right)^{m-j} P_+ \right] - \right. \\ \left. - \left[\left(\frac{1}{D_0} A \right)^j X, \left(\frac{1}{D_0} A \right)^{m-j} P_+ \right] + \left[\left(\frac{1}{D_0} A \right)^j \frac{1}{D_0} [D_0, X], \left(\frac{1}{D_0} A \right)^{m-j} P_+ \right] \right\}$$

Using a geometric series expansion for $(D_A)^{-1} = (D_0 + A)^{-1}$ we can write

$$(16) \quad -i\nu(X; A) = \text{TR} \{ [(D_A)^{-1} D_0, X (D_A)^{-1} D_0 P_+] - [(D_A)^{-1} X D_0, (D_A)^{-1} D_0 P_+] \}.$$

The symbol of the PSDO on the right-hand side can be expanded in a geometric series using

$$\sigma(D_A^{-1} B)(x, p) = (D_0 - i\gamma^\mu \frac{\partial}{\partial x_\mu} + A(x))^{-1} \sigma(B)(x, p)$$

where B is a PSDO. The nontrivial part of the anomaly arises from the γ_{d+1} term in P_+ (see the discussion below on the real part of the anomaly). We shall denote by ν_γ the sum of γ_{d+1} terms in ν . The trace of the commutators is again a boundary integral in momentum space of terms with symbols decaying like $|p|^{-d}$ as $|p| \rightarrow \infty$.

Expanding (16) as geometric series in momentum space,

$$(17) \quad \nu_\gamma(X; A) = \frac{1}{2(2\pi)^d} \int dp \delta(\Lambda - |p|) \int dx \frac{p_k}{|p|} \sum_{m+r=d-1} \text{tr} \left(\frac{1}{\not{p}} \gamma^{i_1} \dots \frac{1}{\not{p}} \gamma^{i_m} \frac{1}{\not{p}} \gamma^{j_1} \dots \frac{1}{\not{p}} \gamma^{j_r} \gamma_{d+1} \right) \times$$

where

$$a_{i_1 \dots i_m} = (-i\partial_{i_1} + A_{i_1}) \dots (-i\partial_{i_m} + A_{i_m}).$$

Taking into account that the nonzero contribution to the trace of the commutators comes from degree $= -d$ terms in momentum space and using the property (9) of the traces of gamma matrices, we obtain

$$(18) \quad \nu_\gamma(X; A) = \frac{1}{2d(2\pi)^d} \text{vol}(S^{d-1}) (2i)^{d/2} (-1)^{(d-2)/2} \sum_{m+r=d-1} (-1)^m \epsilon_{i_1 \dots i_m k j_1 \dots j_r} \int dx \text{tr} \{ (a_{i_1 \dots i_m} \cdot 1) \partial_k (X a_{j_1 \dots j_r} \cdot 1) - (a_{i_1 \dots i_m} \cdot X) \partial_k (a_{j_1 \dots j_r} \cdot 1) \}$$

where $\text{vol}(S^{d-1}) = 2\pi^n / \Gamma(n)$. The antisymmetrized coefficients $a_m(A) = \epsilon_{i_1 \dots i_m} a_{i_1 \dots i_m}$ are evaluated using $F = -idA + A^2$ and the Bianchi identity $-idF + [A, F] = 0$.

We get

$$a_m(A) \cdot Z = \begin{cases} F^{m/2} Z & \text{when } m \text{ even} \\ F^{(m-1)/2} (-idZ + AZ) & \text{when } m \text{ odd} \end{cases}$$

where Z is an arbitrary operator and the products are exterior products of differential forms.

Inserting this to (18) and dropping integrals of total derivatives one gets

$$\nu_\gamma(X; A) \propto \sum_k \int dx \text{tr} dX F^k A F^{n-k-1}.$$

Using the symmetrized trace

$$\text{Str}(T_1, \dots, T_m) = \frac{1}{m!} \sum_{perm} \text{tr} T_{i_1} \dots T_{i_m}$$

the final formula for ν_γ can be compactly written as

$$(19) \quad \nu_\gamma(X; A) = -\frac{(-i)^n}{(n-1)!(2\pi)^n} \int dx \text{Str} dX F^{n-1} A.$$

In the above formula one has to insert dX , A and $(d-2)/2$ copies of F as arguments.

Summarizing, we get for the γ_{d+1} part of the anomaly,

$$(20) \quad \omega_\gamma(X; A) = -\frac{(-i)^n}{(n-1)!(2\pi)^n} \int_0^1 dt \frac{t-1}{t} \int dx \text{Str} (dX A_t F_t^{(d-2)/2})$$

where $A_t = tA$ and $F_t = -itdA + t^2 A^2$.

The real part of the anomaly

The term in $\nu(A; X)$ due to the unit matrix in $P_+ = \frac{1}{2}(1 + \gamma_{d+1})$ is equal to the gauge variation of $\frac{1}{2}\text{TR} \log(1 + \frac{1}{D_0 + i\epsilon} \mathcal{A})$. We have an exact operator equation

$$(21) \quad \begin{aligned} \mathcal{L}_X \log(1 + \frac{1}{D_0 + i\epsilon} \mathcal{A}) &= [\log(1 + \frac{1}{D_0 + i\epsilon} \mathcal{A}), iX] \\ &+ \sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{k(k+1)} \sum_{j=0}^{k-1} \left[(\frac{1}{D_0 + i\epsilon} \mathcal{A})^j (\frac{1}{D_0 + i\epsilon} dX), (\frac{1}{D_0 + i\epsilon} \mathcal{A})^{k-j} \right]. \end{aligned}$$

In contrast to the chiral case the derivation of this equation does not involve the problem that γ_{d+1} does not anticommute with $\frac{1}{D_0 + i\epsilon}$.

The Λ finite part of the cut-off trace of the right-hand side depends only on the projection π_{-d} (because it is a sum of commutators) of the asymptotic expression to the degree $-d$ part. On the other hand, a gauge variation commutes with the degree operator and therefore

$$\begin{aligned} \text{Re } \omega(A; X) &= \mathcal{L}_X \text{tr}_\Lambda \pi_{-d} \frac{1}{2} \log(1 + \frac{1}{D_0 + i\epsilon} \mathcal{A}) \\ &= \mathcal{L}_X P_d(A) \end{aligned}$$

Clearly $P_d(A)$ is a local differential polynomial in A , of degree d . In the case $d = 2$ it is equal to $\frac{1}{8\pi} \int \text{tr} |A|^2 dx$. This argument is not valid for the imaginary part. We do *not* have an exact operator equation like

$$\mathcal{L}_X \log(1 + \frac{1}{D_0 + i\epsilon} \mathcal{A} P_+) = \sum [R_i, T_i]$$

for some operators R_i, T_i .

Actually, one never needs an explicit expression for $P_d(A)$ since in the modified effective action (which has gauge invariant real part) precisely these terms are subtracted. For those operators for which the logarithms are defined we can write

$$(22) \quad \hat{S}_{eff}(A) = \text{TR} \left[\log(1 + \frac{1}{D_0 + i\epsilon} \mathcal{A} P_+) - \frac{1}{2} \pi_{-d} \log(1 + \frac{1}{D_0 + i\epsilon} \mathcal{A}) \right].$$

This is our final formula for the effective action with gauge invariant real part. There is an alternative way to define the trace without using the cut-off mechanism. Since the operator $\frac{1}{D_0} \mathcal{A}$ is of degree -1 the powers $(\frac{1}{D_0} \mathcal{A})^k$ are trace-class when $k > d$. Therefore the operator $\log(1 + \frac{1}{D_0} \mathcal{A} P_+) - \sum_{k=1}^d \frac{(-1)^{k-1}}{k} (\frac{1}{D_0} \mathcal{A})^k P_+$ has a finite trace without any cut-off. However, we have not performed the anomaly calculation for this form of the effective action although we expect to get an equivalent result.

ACKNOWLEDGEMENTS We want to thank G. Ferretti and S. Rajeev for many discussions on anomalies.

REFERENCES

- [BGV] N. Berline, E. Getzler, and M. Vergne: *Heat kernels and Dirac operators*. Springer-Verlag, Berlin (1992)
- [C] A. Connes: *Noncommutative geometry*. Academic Press (1994)
- [CFNW] M. Cederwall, G. Ferretti, B. Nilsson, A. Westerberg: Schwinger terms and cohomology of pseudodifferential operators. **hep-th/9410016**
- [H] L. Hörmander: *The analysis of linear partial differential operators III*. Springer-Verlag, Berlin (1985)
- [L] E. Langmann: Non-commutative integration calculus. **hep-th/9501092**
- [LM] E. Langmann and J. Mickelsson: $(3+1)$ -dimensional Schwinger terms and non-commutative geometry. *Phys. Lett. B* **338**, p. 241 (1994)
- [M] J. Mickelsson: Wodzicki residue and anomalies of current algebras. **hep-th/9404093**. In: *Integrable Models and Strings*, ed. by A. Alekseev, A. Hietamäki, K. Huitu, A. Morozov, and A. Niemi. *Lecture Notes in Physics* **436**, Springer-Verlag, Berlin (1994).
- [R] R. Rennie: Geometry and topology of chiral anomalies in gauge theories. *Adv. Phys.* **39**, p.617 (1990)
- [TJZW] S.B. Treiman, R. Jackiw, B. Zumino, and E. Witten: *Current algebra and anomalies*. Princeton University Press, Princeton (1985)
- [W] M. Wodzicki: Noncommutative residue. In: *Lecture Notes in Mathematics* **1289**, ed. by Yu. Manin, Springer-Verlag, Berlin (1985)